CUMULATIVE INDEX 2000

Volume 18

March GERIATRIC ANESTHESIA, pages 1–215
June REGIONAL ANESTHESIA, pages 217–493
September PERIOPERATIVE MEDICINE, pages 495–703

December ANESTHESIA AND RENAL CONSIDERATIONS, pages 705–994

Note: Page numbers of article and issue titles are in boldface type

Accounting, in business of perioperative medicine, 679–684

Acetaminophen, as analgesic agent in palliative care, 187

Acid-base and electrolyte changes,

physiologic, in acute and chronic renal failure patients, **809–833** hyperkalemia, 819–821, 822

hypermagnesemia, 828–829 hypernatremia, 817–819 hyperphosphatemia, 829–830 hypocalcemia, 821, 823–826 hypomagnesemia, 826–828 hypomagnesemia, 813–817

metabolic acidosis, 810–813 Acute care medicine, community hospital practice, **539–550**

availability of services, 544 critical care, 543–544

hospital organization involvement, 544–545

intraoperative care, 541

opportunities for anesthesiologists in acute care hospitals, 547

preoperative evaluation

preoperative evaluation clinic, 540-541 recruiting and training of

anesthesiologists, 545 success of perioperative medicine, 545–547

surgical clinical pathways, 542–543 uncertain future for anesthesiology, 549–550

working with hospitalist teams, 547–549

Acute cortical necrosis, in pathogenesis of renal insufficiency, 779 Acute tubular necrosis, in pathogenesis of renal insufficiency, 778–779

prerenal syndrome and, interface between, 741–742

as vasomotor nephropathy, 798–799 Adenosine, in cardiopulmonary

resuscitation in the elderly, 152

Advance care planning, 201
Advanced age, as risk factor for acute renal failure, 707

Advance directives, 164–167, 201

Airway(s), in geriatric patients, and effect on cardiopulmonary resuscitation, 148–149

upper and lower, structural alterations in, in elderly, 48–49

Albumin, in kidney transplant surgery, 935–937

Alfentanil, as opioid analgesic, 261 pharmacology of, in elderly patients, 14–15

Ambulation, with neuraxial labor analgesia, 392-393

Ambulatory surgery, outcomes research in, 639–640

regional anesthesia for, **341–353** advantages of, **341–342** paravertebral block, **347–348**

techniques, 342-347 thoracic paravertebral block, 352-353

spinal anesthesia for, 237 Amides, for local anesthesia, bupivacaine, 228–229

dibucaine, 229–230 etidocaine, 229

levobupivacaine, 230 lidocaine, 226–227 Amides (Continued) mepivacaine, 227-228 prilocaine, 229

ropivacaine, 230

Aminosteroids, for renal preservation, 764 Amiodarone, in cardiopulmonary resuscitation in the elderly, 153 Ammonia toxicity, as complication of

transurethral resection of prostate, 892-893

Analgesia, balanced, 585 labor. See Labor analgesia. pre-emptive, 585

from regional anesthesia, 98

Analgesics, nonopioid, for postoperative pain management, in pediatric patients, 622-623

non-opioid, in palliative care, 187, 189 opioid, in palliative care, 189-194 for postoperative pain management, in pediatric patients, 624-626

Anaritide, for prevention of acute renal failure, 713

Anemia, chronic, preoperative evaluation, for kidney transplant surgery, 925-926 in patients with chronic renal failure, 869

Anesthesiologists, opportunities for, in acute care hospitals, 547

recruiting and training, in community hospital practice, 545

Anesthesiology, common problem and special problem in, 519-520 outcomes in perioperative medicine and,

633-645. See also Perioperative medicine, and anesthesiology

Anesthetic drugs, pharmacology of, in elderly patients. See Pharmacology of anesthetic drugs, in elderly patients. Aneurysm, abdominal aortic, after

extracorporeal shock wave lithotripsy, 963-964

Angiotensin-converting enzyme inhibitors. vasomotor nephropathy from, 804-805

Ankle, nerve blocks at, 333-334 Anterior spinal artery syndrome, as complication of spinal and epidural

anesthesia, 468 Antiarrhythmics, in cardiopulmonary resuscitation in the elderly, 152-153

Anticoagulation, oral, spinal hematoma and, 472-473

spinal anesthesia and, 238-239 Antimicrobial agents, as risk factor for acute renal failure, 708

Antiplatelet medications, spinal hematoma and, 473-474

Anuria, intraoperative, causes of, in kidney transplant surgery, 939-940

Aortic cross-clamping, as risk factor for acute renal failure, 710

Aortic stents, endovascular, for prevention of renal failure, 733-734

Aortic surgery, difficulties with assessing literature on perioperative renal protection during, 725-726

difficulties with intraoperative assessment of renal function during, 720-

intraoperative management of renal function during, 719-737

with endovascular aortic stents, 733-

historical perspective, 719-720 with kidney perfusate, 732-733

with pharmacologic agents, calcium channel blockers, 731 dopamine, 729-731 furosemide, 728-729 mannitol, 726-728

with volume loading, 731-732 renal dysfunction caused by, 723-725

Apnea, as intubation-related complication, in pediatric patients, 616-617

postoperative, in premature infants, 604-

as risk for anesthesia in pediatric patients, 606-607

Arginine vasopressin, for septic shock,

vasoconstrictor response and, 794-795 Arrhythmias, in patients with chronic renal failure, 879

Arterial stiffening, as effect of cardiovascular and autonomic aging, 35-36

Arthritis, affecting positioning of aged patients, 108-109

Artificial nutrition and hydration, in terminally ill, 202-203

Asthma, as risk for anesthesia in pediatric patients, 603-604

Atracurium, pharmacology of, in elderly patients, 23

Atrial natriuretic peptide, for oliguria, 747-748

for renal preservation, 759-761 vasodilator response and, 796

Atrial natriuretic peptide analogues, for prevention of acute renal failure, 713

Atropine, decreased response to, as effect of cardiovascular and autonomic aging, 38

Autonomic and cardiovascular function, in geriatric patient, 31-46

effects of anesthesia and hemodynamic management, 40-43

management of hypotension, 42-43

mechanisms of cardiovascular and autonomic aging, changes in sympathetic nervous system, 38–39

connective tissue stiffening, 35–38 decreased beta-receptor responsiveness, 32–35

decreased response to atropine, 38 Axillary blockade, for upper extremity regional anesthesia, 311–312

Back pain, lower, management of, in geriatric patients, 128–131

Bacteremia, transient, as complication of transurethral resection of prostate, 893 Benzocaine, for local anesthesia, 226 Bereavement, 203–204

Beta-receptor responsiveness, decreased, as effect of cardiovascular aging, 32–35

Bladder, neuroanatomy of, 885–886 perforation of, as complication of transurethral resection of prostate, 893

Bleeding, as complication of transurethral resection of prostate, 894

Blockade of stellate ganglion, for chronic pain management, 442–445

Blood loss, with epidural anesthesia and analgesia, 423

Blood transfusion, preoperative, for patients with chronic renal failure, 874–875

874–875
Bowel, affected by extracorporeal shock wave lithotripsy, 963

Bowel obstruction, in terminally ill patients, 196

Brachial plexus block, in pediatrics, 371–372

Brain damage, as cause of CNS dysfunction after anesthesia, 60 Bronchopulmonary dysplasia, as risk for

anesthesia in pediatric patients, 605–606

Bupivacaine, for epidural anesthesia, 260, 261

for local anesthesia, 228–229 for spinal anesthesia, 235–236, 241 transient neurologic symptoms from, 235–236, 238

Burn surgery, as risk factor for acute renal failure, 710

Calcium channel antagonists, in kidney transplant surgery, 938 Calcium channel blockers, for prevention of renal failure, 713, 731

for renal preservation, 753–757

Cardiac disease, preoperative evaluation, for kidney transplant surgery, 922 Cardiac dysfunction, as risk factor for

acute renal failure, 707

Cardiac stiffening, as effect of cardiovascular and autonomic aging, 37–38

Cardiopulmonary bypass, as risk factor for acute renal failure, 710

vasomotor nephropathy from, 803–804 Cardiopulmonary resuscitation, decision not to initiate, in terminally ill, 202 management of, 143–158

outcome of, in elderly patients, 144-147 patients' expectations, 154

physiologic changes with aging and effects on, 147–153

airway, 148–149 cerebral protection, 153 circulation, 149–150 drug therapy, 150–153

Cardiovascular complications, from kidney transplant surgery, 940 Cardiovascular disease, outcomes research

in, 638–639
in patients with chronic renal failure,
868

Cardiovascular function, autonomic and, in geriatric patient. See Autonomic and cardiovascular function, in geriatric patient.

Case-controlled trials, in outcomes research, 636

Catheters, for combined spinal-epidural technique, risk of penetration from, 276–277

for epidural anesthesia, 252 insertion techniques, 252–254 verification of location, 254–255 Caudal anesthesia, for transurethral

resection of prostate, 888 Caudal block, in pediatrics, 356–359 dosing, 358–359

equipment, 357 risks, complications, contraindications, 359

technique, 357-358

Celiac plexus block, for chronic pain management, 447–452

Central nervous system dysfunction after anesthesia, in geriatric patient, 59–70

clinical presentations, abnormal reflexes, 63 cognitive dysfunction after cardiac

surgery, 64 cognitive dysfunction after noncardiac surgery, 64–65

death, 62-63 delirium, 63-64 Central nervous system dysfunction after anesthesia (Continued)

ISPOCD study, 65

methodologic problems in the detection of cognitive dysfunction, 65

postoperative cognitive dysfunction, 64

stroke, 63

mechanisms of, extensive loss of brain tissue, 60

hypoperfusion, 60-61

hypoxemia, 61 infarction, 60

risk factors, 61-62

spinal cord damage associated with central blocks, 62

prevention of, 66 Cerebral death, from CNS damage during

anesthesia, 62-63

Cerebral protection, in cardiopulmonary resuscitation in the elderly, 153

Cervical facet, median branch blocks, for chronic pain management, 432, 434-435

Cesarean section, combined spinalepidural block for, 269-273

general vs. regional anesthesia for, 395-396

postcesarean analgesia for, 396-397 spinal vs. epidural anesthesia for, 396

Chemotherapeutic-immunosuppressive agents, as risk factor for acute renal failure, 708

Children. See Pediatric patients; Pediatrics. 2-Chloroprocaine, for local anesthesia, 225

Chronic pain management, common nerve blocks in. See Nerve blocks, common, in chronic pain management.

Circulation, in geriatric patients, and effect on cardiopulmonary resuscitation,

Circulatory overload, as complication of transurethral resection of prostate,

Cisatracurium, pharmacology of, in elderly patients, 24

Clonidine, in epidural anesthesia, 262 with local anesthetics for lower extremity nerve blocks, 337

for prolonging upper extremity nerve blocks, 306-307

with spinal anesthesia, 246

Coagulation, neuraxial labor analgesia and, 393-394

Coagulopathy, as complication of transurethral resection of prostate, 894 correction of, for patients with chronic renal failure, 875

uremic, preoperative evaluation, for kidney transplant surgery, 926-927

Cocaine, for local anesthesia, 226 Codeine, in palliative care, 190

Cognitive dysfunction, from CNS dysfunction after anesthesia, 64-65 Cohort studies, prospective, in outcomes

research, 635

retrospective, in outcomes research, 636 Combined spinal-epidural technique, 267-295

clinical experience with, 268-276 for cesarean section, 269-273 for day case surgery, 269 for general surgery, 268 in labor, 273-276

for orthopedic surgery, 269 controversial aspects of, controlling upper level of block, 277-278

decreased risk of postdural puncture headache, 282-284

epidural test-doses, 284-285 fetal and neonatal effects, 281 pressure changes in epidural and suba-

rachnoid spaces, 279-281 risk of catheter penetration through dural hole, 276-277

risk of meningitis, 281-282 transdural drug transport, 278-279

for labor, 387 special equipment for, 285-287

Complement system, modulation of, for renal preservation, 763-764 Conduction blockade, physiology of,

218-219 Congestive heart failure, preoperative evaluation, for kidney transplant

surgery, 922-923 Connective tissue stiffening, as effect of cardiovascular and autonomic aging, 35-38

Constipation, in terminally ill patients, 194-195

Continuity of care, in perioperative medicine program, 525

Continuous regional anesthesia, for ambulatory surgery, 348-350

Contoured supine position, advantages of, in aged patients, 116-117 Convection, in dialysis, 855

Coronary artery disease, preoperative evaluation, for kidney transplant surgery, 922

Corticosteroid therapy, as risk for anesthesia in pediatric patients, 609

Cough, in terminally ill patients, 197-198 Critical care, in community hospital practice, 543-544

requirements, 529

and perioperative medicine, 527-538 current perspective, 529-530 current state of the specialty, 530-531 demographic changes and practitioner

historical perspective, 527-529 practice models, 533-536 training paradigm, 531-533 in perioperative medicine program, 524-

Croup, postintubation, in pediatric patients, 617

Cryosurgery, as alternative to transurethral resection of prostate, 895

Cyclosporine A, vasomotor nephropathy from, 805

Cystectomy, radical, anesthetic considerations, 902 preoperative considerations, 901-902 radical prostatectomy, nephrectomy,

pheochromocytoma, and laparoscopic procedures, anesthesia for, 899-917

Data-analysis, in outcomes research, 635-637

Databases, in outcomes research, 636-637 Data management, for perioperative medicine practice, 647-661 constructing a perioperative

information system, 649-650 electronic record design and implementation issues, data standards and information coding, 655-656

hardware issues, 658-659

purchase or develop a system, 658 security, 656-658 selecting important data, 656 training and implementation issues,

659-660 goals of data collection, 647-649 care documentation and professional billing, 648

evidence-based medicine, 649 systems for recording and reporting perioperative information, advanced solution, 652-655 intermediate solution, 651-652

simple solution, 650-651 Day case surgery, combined spinalepidural block for, 269

Decision analysis, in business of perioperative medicine, 689-690 Decubiti, as complication of positioning

aged patients, 113-114 Delirium, from CNS dysfunction after

anesthesia, 63-64 in terminally ill patients, 200

Dementia, in terminally ill patients, 199-200

Depression, in terminally ill patients, 198-199

Diabetes, complications from, in kidney transplant surgery, 941-942 preoperative evaluation, for kidney transplant surgery, 923-924

Dialysis, 853-862

acute and chronic fluid shifts, 859-860 basic mechanisms of, 853, 855

convection, 855 diffusion, 855 osmosis, 855

for chronic renal failure, 874, 880 forms of therapy in, hemodiafiltration,

854, 857 hemodialysis, 855-857 hemofiltration, 854, 857 peritoneal dialysis, 858 ultrafiltration, 854, 857-858

intermittent vs. continuous therapy, 858-

for renal insufficiency, 784-785 surgery in patients on, 860-861 Diarrhea, in terminally ill patients, 196 Dibucaine, for local anesthesia, 229-230 Diffusion, in dialysis, 855

Diltiazem, for renal preservation, 756-757 Diuretic resistance, strategies for overcoming, 749-750 types of, 748-749

Diuretics, classification of, 745 in kidney transplant surgery, 937-938 loop, for oliguria, 746 for renal insufficiency, 783

for oliguria, atrial natriuretic peptide, 747-748

dopamine, 746-747 dopaminergic agonists, 746 fenoldopam, 747 loop diuretics, 746 mannitol, 744, 746

renal benefits, 744 Dobutamine, dopamine used with, 846 Do-not-resuscitate orders in operating room, for geriatric patients, 167-171

Dopamine, for intraoperative prevention of renal failure, 729-731

in kidney transplant surgery, 938–939 low-dose "renal," 835-851 animal studies, 841

in combination therapy, 846 in crossover trial, 846

equivocal evidence, 841, 845 for established acute renal failure,

845-846 history, 836

and oliguric vs. nonoliguric renal failure, 841

in patients at risk for acute renal failure, 845 pharmacodynamics, 837-840

pharmacokinetics, 836-837

Dopamine (Continued)
positive evidence for, 841, 842–844
rationale for use, 840
side effects, 840–841
statistical power of renal dopamine
studies, 845

in management of renal insufficiency, 783

for oliguria, 746-747

for prevention of acute renal failure, 712

Dopamine₁ agonists, for renal preservation, 761–763

Dopaminergic agonists, for oliguria, 746 Dopexamine, in kidney transplant surgery, 939

Dysphagia, from intrathecal analgesia, 388–389

Dyspnea, in terminally ill patients, 196-197

Edema, pulmonary, in postoperative pediatric patients, 618–619

Edmonton symptom assessment scale, 186 Electrolyte balance, fluid and, in patients with chronic renal failure, 864–868

Electrolyte changes, physiologic acid base and. See Acid-base and electrolyte changes, physiologic, in acute and chronic renal failure patients.

Emergence phenomena, after general anesthesia, in pediatric patients, 615–616

End of life issues, anesthesiologist's involvement in, 172–175

palliative care and. See Palliative care, and end of life issues.

Endovascular aortic stents, for prevention of renal failure, 733–734

End-stage liver disease, in pathogenesis of renal insufficiency, 779–780

Epidural abscess, after epidural anesthesia, 477–478, 480

Epidural analgesia, for labor, epidural test dose, 391–392

individualized approach to, 390–391 low dose techniques and patientcontrolled epidural analgesia, 389–390

in perioperative pain management, 581–582

complications, 583–584 controversial issues, 584

potential benefits, 582–583 Epidural anesthesia, and analgesia, new developments in, **251–266** anesthetic and analgesic drugs,

259–263 liposome-encapsulated analgesics, local anesthetic drugs, 260–261 novel analgesic drugs, 261–262 opioid analgesics, 261

equipment, catheters, 252 needles, 251–252

methods of determining extent of epidural anesthesia, 257–258

patient-controlled epidural analgesia, 258–259

techniques, epidural radiographs and epiduroscopy, 256-257

epidural saline injection to hasten return of postoperative motor function, 255–256

needle and catheter insertion techniques, 252–254

unilateral epidural analgesia techniques, 255

verification of catheter location, 254–255

role of, in postoperative outcome, 407-428

benefits of IV-PCA, 409 blood loss and thromboembolism, 423

defining surgical outcome, 410 epidural issues, 409–410 gastrointestinal motility, 424–426 immune system, 423–424

neuroendocrine stress response, 421–423

pain relief, 408–409 physiologic sequelae of surgery,

pulmonary outcome, 418–421 regional vs. general anesthesia in vascular surgery, 410–413

thoracic epidural anesthesia and myocardial ischemia, 414–418

complications of, 461-485

anterior spinal artery syndrome, 468 incidence and etiology of, 462, 463 local anesthetic toxicity, 465–466 meningitis and spinal abscess, 474–

478, 479-480

nerve injury from needle and catheter placement, 462, 464

neuraxial block in patients with preexisting neurologic disorders, 480– 481

performance of neuraxial block in patients under anesthesia, 481–482 postdural puncture headache, 464–465 spinal hematoma, 468

antiplatelet medications, 473–474 intravenous and subcutaneous standard heparin, 470–471

low molecular weight heparin, 471–472

oral anticoagulants, 472-473

transient neurologic symptoms, 466–468

herpes simplex virus and, 478–479 HIV and, 479

vs. spinal anesthesia, for cesarean section, 396

Epidural space, pressure changes in, from combined spinal-epidural technique, 279-281

Epidural steroid injection, for chronic pain management, 438–439

Epidural test-doses, with combined spinalepidural technique, 284–285

Epiduroscopy, 256-257

Epinephrine, in cardiopulmonary resuscitation in the elderly, 150–151

with local anesthetics for lower extremity nerve blocks, 337

for prolonging upper extremity nerve blocks, 306

with spinal anesthesia, 245

Esters, for local anesthesia, benzocaine, 226 2-chloroprocaine, 225 cocaine, 226 procaine, 224

tetracaine, 225–226

Ethical decisions in perioperative elder care, 159–181

advance directives, 164-167

anesthesiologist's involvement in end-oflife care, 172–175

anesthesiologist's stake in medical ethics, 176-177

do-not-resuscitate orders in operating room, 167–171

ethical principles and the concept of personhood, 160-162

informed consent, 162–164

resource allocation, 176 treatment redirection and palliative care, 171–172

views of aging, 159–160

Etidocaine, for local anesthesia, 229 Euthanasia, 172, 174, 175

Extracorporeal shock wave lithotripsy, anesthetic considerations for, in children, 961

immersion physiology and concerns, 960-961

pain, 956-960

complications and special considerations, 961-964

bowel and spleen, 963

heart, 963

hypertension, 962–963 kidney, 962

vascular, 963–964

fundamental lithotriper concepts, 954, 955–956

and percutaneous nephrolithotomy, 953-971 Facet blocks, for chronic pain management, 430–431

Fasting guidelines, preoperative, for pediatric patients, 612–613

Felodipine, for renal preservation, 755–756 Femoral nerve block, 326, 328

complications, 329 in pediatrics, 372

technique, 328-329

Fenoldopam, for oliguria, 747 for renal preservation, 761–763

Fentanyl, as opioid analgesic, 261 in palliative care, 191–192 pharmacology of, in elderly patients,

with spinal anesthesia, 236, 245–246 Fetal effects, from combined spinal-

epidural technique, 281 Fetal heart rate effects, of intrathecal

analgesia, 387–388 Finance, in business of perioperative medicine, 691–692

Fluid and electrolyte balance, in patients with chronic renal failure, 864–868

Fluid imbalance, in patients with chronic renal failure, 864

Fluid resuscitation, impact of, on renal outcome, 741–742

Furosemide, dopamine used with, 846 for prevention of renal failure, 712, 728– 729

Gas exchange, impaired efficiency of, in elderly, 52–53

Gastric aspiration prophylaxis, for kidney transplant surgery, 928

Gastrointestinal dysfunction, in patients with chronic renal failure, 870 Gastrointestinal motility, with epidural

anesthesia and analgesia, 424–426 Gastrointestinal symptoms, in terminally ill patients, 194–196

General anesthesia, improvements in, 99-100

for patients with chronic renal failure, 877–879

spinal anesthesia vs., 236

vs. regional anesthesia, for cesarean section, 395–396

General surgery, combined spinal-epidural block for, 268

General vs. regional anesthesia, choosing, for the elderly, 91–104

anesthesia-related complications, 95–96, 97–98

difference between the late 1970s and the late 1990s, 96–101 early perceptions, 94 General vs. regional anesthesia (Continued)
Hierarchy of Questions about, 91–92
perioperative mortality and
morbidity in older patients,
93–94

recent studies, 96

Geriatric anesthesia, autonomic and cardiovascular function in the geriatric patient, 31–46

central nervous system dysfunction after anesthesia in the geriatric patient, 59–70

choosing general vs. regional anesthesia for the elderly, 91–104

ethical decisions in perioperative elder care, 159–181

geriatric pain management, 123-141 issue on, 1-215

management of cardiopulmonary resuscitation, 143–158

palliative care and end of life issues, 183-208

pharmacology of anesthetic drugs in elderly patients, 1–29 positioning aged patients, 105–121

preoperative assessment of the elderly patient, 71–89

respiratory function in the elderly, 47-58 Glycine toxicity, as complication of transurethral resection of prostate, 892

Grief, stages of, 203–204 Growth factors, for prevention of acute renal failure, 713

for renal repair and regeneration, 765

Head-down tilt, disadvantages of, in aged patients, 115–116

Heart, affected by extracorporeal shock wave lithotripsy, 963

Hematologic alterations, in patients with chronic renal failure, 869

Hemodiafiltration, in dialysis, 854, 857 Hemodialysis procedure, 855–857 Hemodynamic management, effects of

Hemodynamic management, effects of anesthesia and, in geriatric patient, 40-41

Hemofiltration, in dialysis, 854, 857 Hemoglobin, as risk factor for acute renal failure, 709

Heparin, intravenous and subcutaneous, spinal hematoma and, 470–471 low molecular weight, spinal hematoma

and, 471–472 Hepatorenal syndrome, as vasomotor nephropathy, 802–803

Herpes simplex virus, spinal and epidural anesthesia and, 478–479

Herpes zoster, postherpetic neuralgia from, 132–135

HIV, spinal and epidural anesthesia and, 479

Hospice care, 185

Hospitalist teams, in community hospital practice, 547–549

Hydrocodone, in palliative care, 190

Hydromorphone, in palliative care, 191 Hyperkalemia, diagnosis, 821

pathophysiology, 819–820 in patients with chronic renal failure, 879

treatment, 821, 876

Hypermagnesemia, diagnosis, 829 pathophysiology, 828–829 treatment, 829

Hypernatremia, diagnosis, 818 pathophysiology, 817–818

treatment, 818–819 Hyperphosphatemia, diagnosis, 829 pathophysiology, 829

treatment, 829–830 Hypertension, after extracorporeal shock wave lithotripsy in, 962–963

as complication of kidney transplant surgery, 940

control of, for patients with chronic renal failure, 875–876

intraoperative, in patients with chronic renal failure, 879

preoperative evaluation, for kidney transplant surgery, 923

Hypnotics, pharmacology of, in elderly patients, 5–14

Hypocalcemia, diagnosis, 823–824 pathophysiology, 821, 823 treatment, 824–826

Hypomagnesemia, diagnosis, 827 pathophysiology, 826–827 treatment, 827–828

Hyponatremia, clinical features, 813 as complication of transurethral resection of prostate, 891–892

diagnosis, 814–815 pathophysiology, 813 treatment, 815–817

Hypo-osmolality, as complication of transurethral resection of prostate, 891–892

Hypoperfusion, as cause of CNS dysfunction after anesthesia, 60–61 as complication of positioning aged patients, 114–115

Hypotension, as complication of kidney transplant surgery, 941

management of, in geriatric patient, 42-43

maternal, and regional anesthesia for obstetrics, 397-400

Hypothermia, as anesthesia-related complication, 95

as complication of positioning aged patients, 113

as complication of transurethral resection of prostate, 894

Hypovolemia, oliguria and, 739-740 as risk factor for acute renal failure, 707 Hypoxemia, as cause of CNS dysfunction after anesthesia, 61

Hypoxia, medullary, in pathogenesis of renal insufficiency, 776-778

Ileus, postoperative, 576-577 Ilioinguinal and iliohypogastric nerve block, in pediatrics, dosing, 367 risks, complications, contraindications, 368 technique, 367

Immune system, preservation of, with epidural anesthesia and analgesia, 423-424

Immunosuppressants, anesthetic implications of, in kidney transplant surgery, 943, 944

Induction agents, intravenous, for kidney transplant surgery, 929

Infarction, as cause of CNS dysfunction after anesthesia, 60

Information management system, perioperative, 520-521

Informed consent, in elderly population, 162-164

Infraclavicular block, for upper extremity regional anesthesia, 310-311

Inguinal perivascular block, for lumbar plexus block, 320-321, 322

Inhalational agents, for kidney transplant surgery, 929-930

Interscalene block, for upper extremity regional anesthesia, 308-309

Intracardiac shunts, as risk for anesthesia in pediatric patients, 607 Intracranial pathology, affecting

positioning of aged patients, 112-113 Intraoperative care, in community hospital practice, 541

in perioperative medicine program, 523 Intrathecal adjuncts, with spinal anesthesia, 236, 245-246

Intrathecal analgesia, for labor, 384-387 adverse effects of, fetal heart rate effects, 387-388

high block, respiratory depression, and dysphagia, 388-389 pruritus, 389

Intravascular volume, in kidney transplant surgery, 933-935

Intravenous regional anesthesia, for upper extremity nerve blocks, 312-313

Intubation-related complications, in pediatric patients, 616-617

Irrigation solutions, absorption of, as complication of transurethral resection of prostate, 891

for transurethral resection of prostate, 884-885

Isolated peripheral nerve blocks, for upper extremity regional anesthesia, 313-314

Jaundice, cholestatic, as risk factor for acute renal failure, 707 postoperative, in pediatric patients, 626-

Juxtaglomerular apparatus, in normal regulation of renal perfusion, 792

Ketamine, in epidural anesthesia, 262 Kidney, conditions involving, after extracorporeal shock wave lithotripsy,

harvested, preservation of, 933 Kidney cocktail, in kidney transplant surgery, 939

Kidney donors, cadaveric, management of, 932-933

living, management of anesthesia for,

Kidney perfusate, for intraoperative prevention of renal failure, 732-733

Kidney recipients, anesthetic management of, 933

Kidney transplant surgery, anesthesia for, 919-951

anesthetic implications of immunosuppressants, 943, 944 candidates for, 919-920

contraindications to, 919, 920 factors affecting viability in, albumin, 935-937

anesthetic management of kidney recipients, 933

calcium channel antagonists, 938 diuretics, 937-938

dopamine, 938-939

dopexamine, 939

intravascular volume, 933-935 kidney cocktail, 939

management of anesthesia for living kidney donors, 932

management of cadaveric kidney donor. 932-933

mannitol, 938

mechanical and other causes for intraoperative anuria and oliguria, 939-940

Kidney transplant surgery (Continued)

preservation of harvested organ, 933 intraoperative complications, cardiovascular complications, 940 diabetes and intraoperative course, 941-942 hypertension, 940 hypotension, 941 intraoperative potassium and glucose levels, 941 sickle cell disease, 942 perioperative management, gastric aspiration prophylaxis, 928 inhalational agents, 929-930 intravenous induction agents, 929 monitoring, 928-929 muscle relaxants, 930-931 premedication, 927-928 selection of anesthesia, 927 use of opioids in patients with uremia, 930 postoperative management, emergence and immediate postoperative course, 943 long-term outcome, 945 preoperative evaluation, 920-927 cardiac disease, 922 chronic anemia, 925-926 congestive heart failure, 922-923 coronary artery disease, 922 diabetes mellitus, 923-924 hypertension, 923 serum potassium, 924-925 uremic coagulopathy, 926-927 Knee, nerve blocks at, 331-332 Labor analgesia, adverse effects of, 387-389

ambulating with, 392-393 combined spinal-epidural analgesia for, 273-276 epidural, 389-392 intrathecal analgesia, 384-387 pain pathways in labor and targeted therapy, 383-384 role of combined spinal-epidural technique, 387 Laparoscopic procedures, anesthetic management, analgesia, 912 anesthetic agents, 911-912 anesthetic technique, 911 antiemesis, 912-913 complications, 913 monitoring, 910-911 muscle relaxants, 912 preoperative evaluation, 910 cardiovascular effects, 907-908 radical prostatectomy, cystectomy, nephrectomy, and pheochromocytoma, anesthesia for, 899-917

respiratory effects, 908-910 Laser prostatectomy, as alternative to transurethral resection of prostate, 895 Latex allergy, in pediatric patients, 610-612 "Lawn chair" position, advantages of, in aged patients, 116-117 Leadership, in business of perioperative medicine, 684-685 Levobupivacaine, for epidural anesthesia, 261 for local anesthesia, 230 Levorphanol, in palliative care, 192 Lidocaine, for local anesthesia, 226-227 for spinal anesthesia, 235, 241 transient neurologic symptoms from, 235, 237 Liposome-encapsulated analgesics, in epidural anesthesia, 263 Lithotripsy, extracorporeal shock wave. See Extracorporeal shock wave lithotripsy. Liver disease, end-stage, in pathogenesis of renal insufficiency, 779-780 Liver failure, as vasomotor nephropathy, 802-803 Liver transplantation, as risk factor for acute renal failure, 710 Living wills, 164-165, 201 Local anesthetics, for lower extremity nerve blocks, 319-340 additives, 337 clonidine, 337 epinephrine, 337 sodium bicarbonate, 337 concentration, 336-337 duration, 336 pharmacology of, 217-233 basic nerve cell physiology, 217-218 biochemistry of local anesthetic molecules, 220-223 allergy, 224 cardiac toxicity, 223 direct tissue toxicity, 223-224 lipid solubility, 221 methemoglobinemia, 223 pKa, 221-222 protein binding, 221 toxicity, 222-223 physiology of conduction blockade, 218-219 specific agents, amides, bupivacaine, 228-229 dibucaine, 229-230 etidocaine, 229 levobupivacaine, 230 lidocaine, 226-227 mepivacaine, 227-228 prilocaine, 229 ropivacaine, 230 esters, benzocaine, 226 2-chloroprocaine, 225

cocaine, 226 procaine, 224 tetracaine, 225–226

structure of local anesthetic molecule, aromatic group, 219–220 hydrophilic group, 220 intermediate chain and amide/ester

bond, 220

toxicity from, 222-224

as complication of spinal and epidural anesthesia, 465–466

in pediatrics, 373–376 Loop diuretics, for oliguria, 746 for renal insufficiency, 783

Lower back pain, management of, in geriatric patients, 128–131

Lower extremity nerve blocks, 319–340 femoral nerve block, 326, 328

complications, 329 technique, 328-329

local anesthetic agents, additives, clonidine, 337

epinephrine, 337 sodium bicarbonate, 337 concentration, 336–337

concentration, 336-337 duration, 336

lumbar plexus block, complications, 323 inguinal perivascular block, 320–321, 322

psoas compartment block, 321–323, 324

nerve blocks at the ankle, 333–334 complications, 336

technique, 335 nerve blocks at the knee, 331–332 complications, 333

technique, 332 obturator nerve block, 329–330, 331 complications, 330 technique, 330

sciatic nerve block, 235, 323 anterior technique, 325–326, 327 classic technique, 325 complications, 326

Lower respiratory disease, postoperative, in pediatric patients, 618–619

Lumbar epidural, in pediatrics, 359–364 dosing, 361–362 equipment, 360–361

risks, complications, contraindications, 362–363

technique, 361 test dose, 363-364

Lumbar facet, median branch blocks, for chronic pain management, 431–432

Lumbar neuraxial blocks, regional anesthetic alternatives to, 394–395 Lumbar plexus block, with inguinal

perivascular block, 320–321, 322 with psoas compartment block, 321–323, 324 Lumbar sympathetic blockade, for chronic pain management, 445–447 Lung volumes, changes in, in elderly,

49-52

Magnesium, in cardiopulmonary resuscitation in the elderly, 151–152 and potassium balance, in patients with chronic renal failure, 865–867

Managerial effectiveness, in business of perioperative medicine, 687–689

Mannitol, for intraoperative prevention of renal failure, 726–728 in kidney transplant surgery, 938 for oliguria, 744, 746

for prevention of acute renal failure, 712 Marketing, in business of perioperative

medicine, 690–691 Maternal hypotension, and regional anesthesia for obstetrics, 397–400

Mechanical ventilation, discontinuing, in terminally ill, 203

Medicare Hospice Benefit, 185

Medullary hypoxia, in pathogenesis of renal insufficiency, 776–778

α-Melanocyte-stimulating hormone, for renal preservation, 764–765

Memorial Symptom Assessment Scale, 187, 188–189

Meningitis, as complication of spinal and epidural anesthesia, 474, 476–477, 479–480

risk of, from combined spinal-epidural technique, 281–282

Meperidine, in palliative care, 192 for spinal anesthesia, 242

Mepivacaine, for local anesthesia, 227–228 for spinal anesthesia, 235, 242 transient neurologic symptoms from,

235, 237–238 Meta-analysis, in outcomes research, 637–638

Metabolic acidosis, diagnosis, 813 intraoperative management, 813 pathophysiology, 810–812 in patients with chronic renal failure, 864–865

treatment, 813

Metabolic and nutritional problems, in patients with chronic renal failure, 869–870

Methadone, in palliative care, 192 Methemoglobinemia, from local anesthetics, 223

Microeconomics, in business of perioperative medicine, 690

Microwave ablations, as alternative to transurethral resection of prostate, 895 Midazolam, pharmacology of, in elderly patients, 12–14

Morphine, in palliative care, 191 Murmurs, as risk for anesthesia in pediatric patients, 607–608

Muscle relaxants, for kidney transplant surgery, 930–931

for laparoscopic procedures, 912 for patients with chronic renal failure, 878

pharmacology of, in elderly patients, 23–26

Myalgia, succinylcholine-induced, in postoperative pediatric patients, 626

Myocardial ischemia, thoracic epidural anesthesia and, 414–418

Myoglobin, as risk factor for acute renal failure, 709

Nausea, postoperative, 577 in pediatric patients, 619–621 and vomiting, in terminally ill patients, 195–196

Needles, for combined spinal-epidural technique, 285–287

for epidural anesthesia, 251–252 insertion techniques, 252–254

for preventing post-dural puncture headache, 239–240

Negotiation, in business of perioperative medicine, 695–697

Neonatal effects, from combined spinalepidural technique, 281

Neostigmine, with spinal anesthesia, 246 Nephrectomy, radical, anesthetic considerations, 903

preoperative considerations, 902–903 radical prostatectomy, cystectomy, pheochromocytoma, and laparoscopic procedures, anesthesia for, 899–917

Nephrolithotomy, percutaneous. See Percutaneous nephrolithotomy.

Nephropathy, toxic, in pathogenesis of renal insufficiency, 780–781

vasomotor. See Vasomotor nephropathy. Nephrotoxin exposure, oliguria and, 740 as risk factor for acute renal failure, 708– 709

vasomotor nephropathy from, 804-805 Nerve blocks, at the ankle, 333-334

complications, 336 technique, 335

common, in chronic pain management, 429–459

somatic nerve blocks, blockade of stellate ganglion, 442–445 celiac plexus block, 447–452 cervical facet, median branch blocks, 432, 434–435 epidural steroid injection, 438-439 facet blocks, 430-431

lumbar facet, median branch blocks, 431-432

lumbar sympathetic blockade, 445–447

sacroiliac joint injection, 435–438 selective nerve root block, 439– 441

superior hypogastric plexus block, 452–454

sympathetic nerve blocks, 442 at the knee, 331–332

complications, 333 technique, 332

lower extremity. See Lower extremity nerve blocks.

peripheral, in perioperative pain management, 580–581 upper extremity. See *Upper extremity*

nerve blocks.

Nerve cell physiology, and local anesthetics, 217–218

Neuraxial blockade, in patients under anesthesia, 481–482

in patients with preexisting neurologic disorders, 480–481

Neuraxial labor analgesia ambulating with, 392–393 and coagulation, 393–394

lumbar, regional anesthetic alternatives to, 394–395

Neuroendocrine stress response, epidural anesthesia and analgesia and, 421–423 Neurologic complications, in patients with

chronic renal failure, 870–871 Neurologic injury, from spinal anesthesia, 238–239

Neuromuscular diseases, as risk for anesthesia in pediatric patients, 608–609

Neuropathies, as complication of positioning aged patients, 117–118

Neuropathy, pain management of, in geriatric patients, 131–132

Nitric oxide, role of, in vasodilator response, 797–798

Nonopioid analgesics, in palliative care, 187, 189

for postoperative pain management, in pediatric patients, 622–623

Nonsteroidal anti-inflammatory drugs (NSAIDs), as analgesic agent in palliative care, 187, 189

as risk factor for acute renal failure, 708-

vasomotor nephropathy from, 804 Norepinephrine, dopamine used with, 846 for septic shock, 801 Nutritional and metabolic problems, in patients with chronic renal failure, 869–870

Obesity, affecting positioning of aged patients, 106–108

Obstetric anesthesia, outcomes research in, 640

Obstetrics, regional anesthesia for, 383–406 adverse effects of intrathecal analgesia, fetal heart rate effects, 387–388

> high block, respiratory depression, and dysphagia, 388–389

pruritus and other effects, 389 alternatives to lumbar neuraxial blocks, 394–395

ambulating and neuraxial labor analgesia, 392–394 neuraxial blocks and coagulation,

393–394 cesarean section, general vs. regional

anesthesia, 395–396 epidural analgesia, epidural test dose,

391–392 individualized approach to labor

analgesia, 390–391 low dose techniques and patientcontrolled epidural anesthesia, 389–390

labor analgesia, intrathecal analgesia, 384–387

pain pathways in labor and targeted therapy, 383–384

role of combined spinal-epidural technique in, 387

postcesarean analgesia, 396–397 ropivacaine, 400 spinal vs. epidural anesthesia, 396

uterine perfusion, maternal hypotension, vasopressors, 397–400

Obstructive sleep apnea, as risk for anesthesia in pediatric patients, 606–607

Obturator nerve block, 329–330, 331 complications, 330 technique, 330

Oliguria, clinical approach to, general principles, 742–744

in the ICU, systematic approach to diagnosis and treatment, 739–752

intraoperative, causes of, in kidney transplant surgery, 939–940 pharmacology of diuretic agents for,

744–750 and prerenal syndrome, 739–742

Operations management, in business of perioperative medicine, 693–695

Opioids, in palliative care, 189–194 in patients with uremia, 930 pharmacology of, in elderly patients, 14–23

for postoperative pain management, in pediatric patients, 624–626

for prolonging upper extremity nerve blocks, 307

for spinal anesthesia, 242

systemic, in perioperative pain management, 580

Organizational design, in business of perioperative medicine, 693

Orthopedic surgery, combined spinalepidural block for, 269

Osmosis, in dialysis, 855

Osteoarthritis, pain management for, in geriatric patients, 126–127

Osteomalacia, affecting positioning of aged patients, 108

Osteoporosis, affecting positioning of aged patients, 108

Outcomes research, in anesthesiology, future directions, evidence-based approach to decision making and medical technology assessment, 642–643

expansion into new and nontraditional areas, 642 movement toward nontraditional

outcomes, 641–642 current status of, in perioperative medicine and anesthesiology, selected topics, ambulatory surgery, 639–640

> cardiovascular disease, 638–639 obstetric and regional anesthesia, 640

pain management, 640–641 evolution of, in medicine, 633–634 methodologic issues in, data types and data analysis, meta-analysis, 637–

prospective data, 635–636 retrospective data, 636–637 definition of outcomes research, 634

use of surrogate endpoints, 634–635 Oxycodone, in palliative care, 190–191

Pain, assessing symptoms of, in palliative care, 187

Pain management, chronic. See Chronic pain management.

geriatric, anesthesiologist's perspective, 123–141

geriatric pain, 124–125 for lower back pain, 128–131 for neuropathy, 131–132 Pain management (Continued) for osteoarthritis, 126-127 physiologic effects of aging, 123-124 for postherpetic neuralgia, 132-135 outcomes research in, 640-641 in palliative care, with non-opioid analgesic agents, 187, 189 with opioid analgesic agents, 189-194 perioperative. See Perioperative pain management. Palliative care, clinical skills for, 184 death in the United States, 184-185 definition of, 183-184 end-of-life, anesthesiologist's involvement in, 172-175 and end of life issues, 183-208 advance care planning, 201 bereavement, 203-204 breaking bad news, 201-202 communication and the physician's role, 200-201 life sustaining treatment in the terminally ill, 202-203 opportunities for growth near the end of life, 204 for geriatric patients, 171-172 Medicare Hospice Benefit, 185 prognostication in, 184 symptoms and their treatment, formal assessment, 185-187 gastrointestinal symptoms, bowel obstruction, 196 constipation, 194-195 diarrhea, 196 nausea and vomiting, 195-196

non-opioid analgesic agents, 187, 189 opioid analgesic agents, 189–194 pain symptoms, 187 psychiatric symptoms, delirium, 200 dementia, 199–200 depression, 198–199 respiratory symptoms, cough, 197–198 dyspnea, 196–197 Pancuronium, pharmacology of, in elderly patients, 23

Paravertebral blocks, for ambulatory anesthesia, 347–348 Patient-controlled analgesia, benefits of, 409

Patient-controlled epidural analgesia, 258–259

Pediatric patients, extracorporeal shock wave lithotripsy in, 961

perioperative management issues in. See Perioperative management issues, in pediatric patients.

Pediatrics, regional anesthesia in, 355–381, 481–482 brachial plexus block, 371–372 caudal block, 356–359 dosing, 358–359
equipment, 357
risks, complications,
contraindications, 359
technique, 357–358
femoral nerve block, 372
ilioinguinal and iliohypogastric nerve
block, dosing, 367
risks, complications,
contraindications, 368
technique, 367

technique, 367 local anesthetic toxicity, 373–376 lumbar epidural, 359–364 dosing, 361–362 equipment, 360–361 risks, complications, contraindications, 362–363 technique, 361 test dose, 363–364 penile nerve block, 368–371

dosage, 369
risks and complications, 370–371
technique of Bacon, 369
technique of Dalens, 369
technique of Yaster and Maxwell,
370

peripheral nerve blockade, 366 ring block of the penis, use of nerve stimulator, 371 ropivacaine, 376 sciatic nerve block, 372–373 spinal subarachnoid anesthesia, 364–366

dosing, 365–366 risks and complications, 366 technique, 365

Penile nerve block, in pediatrics, 368–371 dosage, 369 risks and complications, 370–371 technique of Bacon, 369 technique of Dalens, 369 technique of Yaster and Maxwell, 370 Penis, neuroanatomy of, 887

Percutaneous nephrolithotomy, anesthetic considerations for, 964–965 complications and special considerations, 965–966

extracorporeal shock wave lithotripsy and, 953–971 fundamental concepts of, 964

Perioperative information management system, 520–521 Perioperative management issues, in

pediatric patients, 601–632 fasting guidelines, 612–613 latex allergy, 610–612 postoperative care issues, early postoperative anesthesic problems, 613–614 emergence phenomena following

general anesthesia, 615–616

intubation-related complications, 616-617 jaundice, 626-627 lower respiratory disease, 618-619 nausea and vomiting, 619-621 pain management, nonopioid analgesics, 622-623 opioids, 624-626 respiratory problems following general anesthesia, 616 succinylcholine myalgia, 626 upper airway obstruction, 617-618 preoperative pregnancy testing, 610 risks of anesthesia, cardiovascular, intracardiac shunts, 607 murmurs, 607-608 subacute bacterial endocarditis prophylaxis, 608 chronic steroid therapy, 609 neuromuscular diseases, 608-609 respiratory, asthma, 603-604 bronchopulmonary dysplasia, 605-606 obstructive sleep apnea, 606-607 prematurity and apnea, 604-605 pulmonary function tests, 604 upper and lower respiratory tract infections, 602-603 sickle cell anemia, 609-610 Perioperative medicine, and anesthesiology, current status of outcomes research in, ambulatory surgery, 639-640 cardiovascular disease, 638-639 obstetric and regional anesthesia, pain management, 640-641 evolution of outcomes research in medicine, 633-634 future directions, evidence-based approach to decision making and medical technology assessment, 642-643 expansion into new and nontraditional areas, 642 movement toward nontraditional outcomes, 641-642 methodologic issues in outcomes research, data types and dataanalysis, 635-638 definition of outcomes research, 634 use of surrogate endpoints, 634-635 outcomes in, 633-645 business of, 677-698 accounting, 679-684 decision analysis, 689-690 finance, 691-692

leadership, 684-685

marketing, 690-691

managerial effectiveness, 687-689

microeconomics, 690 negotiation, 695-697 operations management, 693-695 organizational design, 693 power, politics, and influencing others. 685-687 strategy, 692-693 critical care and, 527-538 current perspective, 529-530 current state of the specialty, 530-531 demographic changes and practitioner requirements, 529 historical perspective, 527-529 practice models, 533-536 training paradigm, 531-533 data management for practice in, 647-661 constructing a perioperative information system, 649-650 electronic record design and implementation issues, data standards and information coding, 655-656 hardware issues, 658-659 purchase or develop a system, 658 security, 656-658 selecting important data, 656 training and implementation issues, 659-660 goals of data collection, 647-649 care documentation and professional billing, 648 evidence-based medicine, 649 systems for recording and reporting perioperative information, advanced solution, 652-655 intermediate solution, 651-652 simple solution, 650-651 experiment in, 517-519 good results, 518-519 problems, 519 future of anesthesiology is, 495-513 clinical practice of perioperative medicine, 497-498 evolution of a new paradigm for anesthesiology, 496-497 responding to threats to anesthesiology, boundaries of new specialty, 507-508

change management, 505–506 changes in health care environment, 499–500 changes in practice patterns, 502

changes in residency training, 506–507 economic factors, 504–505

implementation of perioperative medicine program, 504 informatics, 508–509 lack of understanding regarding anesthesiology services, 498–499 Perioperative medicine (Continued) role of nonphysicians, 507 third path, 509-510 threats to research, 502-503 workforce issues, 500-502 ideal concept of, 516-517 issue on, 495-703 success of, in community hospital practice, 545-547 Perioperative pain management, current trends in, 575-599 evolution from postoperative pain control to multimodal postoperative care and rehabilitation, 584-586 future, 590-591 options, epidural analgesia, 581-584 peripheral nerve blocks, 580-581 systemic opioids, 580 pathophysiology of postoperative period, coagulation system, cognitive dysfunction, 578 endocrine and metabolic stress response, 575-576 gastrointestinal effects, 576-577 immunologic response, 577 pulmonary function, 576 practical concerns regarding development of, 586, 588-590 financial considerations, 588-589 physician education, training, and certification, 589-590 potential added value of pain management, 590 staffing, 589 rationale for, 578-580 Perioperative physicians, training, 515-525 program for, continuity of care, 525 critical care medicine, 524-525 intraoperative management, 523 post anesthesia acute care, 524 postoperative ward medicine, 524 preoperative evaluation, testing and risk assessment, 522-523 Peripheral airway and lung disease, postoperative, in pediatric patients, Peripheral nerve blocks, in pediatrics, 366 in perioperative pain management, 580-581 Peritoneal dialysis, 858 Pharmacologic options, new, for renal

preservation, 753-771

atrial natriuretic peptide, 759-761 calcium channel blockers, 753-757

α-melanocyte-stimulating hormone,

dopamine₁ agonists, 761-763

aminosteroids, 764

764-765

modulation of the complement system, 763-764 prostaglandins, 757-759 renal repair and regeneration, 765 Pharmacology of anesthetic drugs, in elderly patients, 1-29 effects of aging on, 1-5 body composition, 2-3 cardiovascular effects, 3 end-organ sensitivity, 5 hepatic effects, 3 protein binding effects, 4-5 renal effects, 3-4 variability, 5 hypnotics, 5-14 midazolam, 12-14 propofol, 6-7, 9-12 thiopental, 5-6, 7, 8 muscle relaxants, 23-26 atracurium, 23 cisatracurium, 24 pancuronium, 23 vecuronium, 24-26 opioids, alfentanil, 14-15 fentanyl, 14-15 remifentanil, 16-23 sufentanil, 15-16 Phenylephrine, dopamine used with, 846 Pheochromocytoma, anesthetic considerations, 905-907 preoperative considerations, 904-905 radical prostatectomy, cystectomy, nephrectomy, and laparoscopic procedures, anesthesia for, 899-917 Phosphate, calcium, and renal osteodystrophy, in patients with chronic renal failure, 867-868 Physician-assisted suicide (PAS), 172, 174-175 Positioning aged patients, 105-121 geriatric characteristics, body habitus, 106-108 intracranial pathology, 112-113 skeleton, 108-109 vasculature, 109-112 patient movement, 119-120 potential complications, hypoperfusion, 114-115 hypothermia, 113 neuropathies, 117-118 posture and treatment of hypotension, 115-117 preanesthetic evaluation, 118-119 skin damage and decubiti, 113-114 Post anesthesia acute care, in perioperative medicine program, 524 Postdural puncture headache, choice of needle and, 239-240 as complication of spinal and epidural anesthesia, 464-465

decreased risk of, with combined spinalepidural technique, 282-284

Postherpetic neuralgia, pain management of, in geriatric patients, 132–135 Postintubation croup, in pediatric patients,

617

Postoperative care, in community hospital practice, 542

Postoperative outcome, role of epidural anesthesia and analgesia in. See Epidural anesthesia and analgesia, role of, in postoperative outcome.

Postoperative ward medicine, in perioperative medicine program, 524 Potassium and magnesium balance, in patients with chronic renal failure,

865-867

Power, politics, and influencing others, in business of perioperative medicine, 685–687

Power of attorney for health care (PAHC), 165–166, 201

Pregnancy testing, preoperative, in adolescents, 610

Premature infants, postoperative apnea in, 604–605

Premedication, for kidney transplant surgery, 927–928

for patients with chronic renal failure, 876

Preoperative assessment, of the elderly patient, 71–89

for correct positioning, 118–119 evaluation, 71–80

age-related diseases, 77–78 determining functional reserve, 73–74

drug interactions and altered pharmacology, 78–80 organs at risk, 74–77

cardiopulmonary function, 75–76 hepatorenal and metabolic functions, 76–77

nervous system, 75 physical status and perioperative risk, 83–84

planning an anesthetic, 84–85 preoperative testing, 80–83 in perioperative medicine program, 522–

Preoperative evaluation clinic, in

community hospital practice, 540–541 Preoperative preparation, value,

perspective, and practice in patient care, 551–574

definition of terms, 552–554 future trends, 568–570 perspective, of patient, 556–558 of payor, 562–563 of provider, 558–562 practice, 563-568 value, 554-556

Prerenal syndrome, and acute tubular necrosis, interface between, 741–742 oliguria and, 739–741

Prilocaine, for local anesthesia, 229 Procaine, for local anesthesia, 224

for spinal anesthesia, 241

transient neurologic symptoms from, 235, 238

Propofol, pharmacology of, in elderly patients, 6–7, 9–12

Propoxyphene, in palliative care, 191 Prostaglandins, for renal preservation, 757–759

vasodilator response and, 796

Prostate, neuroanatomy of, 886-887

transurethral resection of, 883–897 advantages of regional anesthesia over general anesthesia, 888–891

alternatives to, 895

anesthetic techniques, 888

complications, absorption of irrigating solution, 891

ammonia toxicity, 892–893 bleeding and coagulopathy, 894 circulatory overload, hyponatremia,

and hypo-osmolality, 891–892 glycine toxicity, 892

hypothermia, 894 perforation of bladder, 893

transient bacteremia and septicemia, 893

TURP syndrome, 894-895 irrigation solutions, 884-885

neuroanatomy of penis, prostate, and bladder, 885–887

pathophysiology of hypertrophic prostate, 883–884

surgical procedures, 884

Prostatectomy, radical, 899–890 anesthetic considerations, 900–901 cystectomy, nephrectomy,

> pheochromocytoma, and laparoscopic procedures, anesthesia for, 899–917

preoperative considerations, 900 Pruritus, from intrathecal analgesia, 389 Psoas compartment block, for lumbar

plexus block, 321–323, 324 Psychiatric symptoms, in terminally ill

patients, 198–200 Pulmonary edema, postoperative, in pediatric patients, 618–619

Pulmonary function, in patients with chronic renal failure, 868

Pulmonary function tests, before anesthesia in pediatric patients, 604 Pulmonary outcome, after epidural

anesthesia and analgesia, 418–421

Radiocontrast media, as risk factor for acute renal failure, 708

Radiographs, epidural, 256-257

Randomized, controlled trial, in outcomes research, 635–636

Reflexes, abnormal, from CNS dysfunction after anesthesia, 63

Regional anesthesia, advantages of, for transurethral resection of prostate, 888–891

for ambulatory surgery, 341-353

analgesia from, 98

combined spinal-epidural technique, 267–295

common nerve blocks in chronic pain management, 429–459

complications of spinal and epidural anesthesia, 461–485

failure rate of, 100-101

hemodynamic control during, 100

issue on, 217-493

lower extremity nerve blocks, 319–340 new developments in epidural anesthesia and analgesia, 251–266

new developments in spinal anesthesia, 235–249

for obstetrics, 383-406

outcomes research in, 640

for patients with chronic renal failure, 877

in pediatrics, 355-381

pharmacology of local anesthetics, 217–233

reasons for safety of, 92-93

role of epidural anesthesia and analgesia in postoperative outcome, 407–428 sedation during, 99

upper extremity nerve blocks, 297–317 volume administration during, 100

vs. general. See General vs. regional anesthesia, choosing, for the elderly.

Remifentanil, pharmacology of, in elderly patients, 16–23

Renal autoregulation, as vasodilator response, 796–797

Renal blood flow, in pathogenesis of renal insufficiency, 776

Renal blood flow regulation, autoregulation, and vasomotor nephropathy, 791–807

Renal considerations, anesthesia and, issue on, 705–994

Renal dysfunction, in aortic surgery patients, caused by surgery, 723–725 preoperative, as risk factor for acute renal failure, 707

Renal failure, acute. See also Renal insufficiency. evolution of care for, 705–706 patient at risk for, 705–717 risk factors for, high-risk surgical procedures, 709–710

nephrotoxins, 708-9 patient-based, 706-707

strategies to prevent, pharmacologic approaches, 711-713

preoperative optimization, 710–711 acute and chronic, physiologic acid-base and electrolyte changes in, **809**–

> hyperkalemia, 819–821, 822 hypermagnesemia, 828–829 hypernatremia, 817–819 hyperphosphatemia, 829–830 hypocalcemia, 821, 823–826 hypomagnesemia, 826–828

hyponatremia, 813–817 metabolic acidosis, 810–813

anesthetic considerations for the patient with, 863-882

pathophysiologic effects of chronic renal failure, cardiovascular disease, 868

fluid and electrolyte balance, 864-868

gastrointestinal dysfunction, 870 hematologic alterations, 869 neurologic complications, 870–871 nutritional and metabolic problems, 869–870

pulmonary function, 868

in aortic surgery patients, pharmacologic agents for intraoperative prevention of, calcium channel blockers, 731

> dopamine, 729–731 furosemide, 728–729 mannitol, 726–728

chronic, pharmacologic alterations in, 871–873

oliguric vs. nonoliguric, dopamine and, 841

perioperative, risk factors for, 740 perioperative management of patient with, general anesthesia, anesthe-

tic emergence, 879 anesthetic induction, 877–878 choice of muscle relaxant, 878

hyperkalemia and arrhythmias, 879 intraoperative hypertension, 879 maintenance of anesthesia, 878–879

intraoperative preparation, 876–877 postoperative care, laboratory investigations, 880

monitoring, 879–880

postoperative fluid management and dialysis, 880

preoperative evaluation, 873 preoperative preparation, blood transfusion, 874–875 control of hypertension, 875–876 correction of coagulopathy, 875 dialysis, 874 premedication, 876

treatment of hyperkalemia, 876 regional anesthesia, 877

Renal function in aortic surgery patients, difficulties with intraoperative assessment of, 720–723

historical perspective, 719-720

intraoperative management of, 719–737 with endovascular aortic stents, 733–

with kidney perfusate, 732–733 with pharmacologic agents, calcium

channel blockers, 731 dopamine, 729–731 furosemide, 728–729

mannitol, 726–728 with volume loading, 731–732

Renal insufficiency, pathophysiology and management of, in perioperative and critically ill patient, 773–789

assessment, 781–782 etiology, 774–775

management, 782-785 pathogenesis, acute cortical necrosis,

acute tubular necrosis, 778–779 end-stage liver disease, 779–780

medullary hypoxia, 776–778 renal blood flow, 776 sepsis, 779

toxic nephropathy, 780-781

Renal perfusion, normal regulation of, 791–798

extrinsic control mechanisms, 792 juxtaglomerular apparatus, 792 renal autoregulation, 796–797 role of nitric oxide, 797–798

vasoconstrictor responses, arginine vasopressin, 794–795 renin-angiotensin-aldosterone axis,

793–794 sympathoadrenal activation, 792–793

vasodilator responses, atrial natriuretic peptide, 796 prostaglandins, 796

Renal preservation, new pharmacologic options for, 753–771

aminosteroids, 764

atrial natriuretic peptide, 759–761 calcium channel blockers, 753–757 dopamine₁ agonists, 761–763 α-melanocyte-stimulating hormone,

764–765 modulation of the complement system, 763–764

prostaglandins, 757-759

renal repair and regeneration, 765 Renal protection in aortic surgery patients, perioperative, difficulties with

assessing literature on, 725–726 Renal repair and regeneration, growth factors for, 765

Renin-angiotensin-aldosterone axis, vasoconstrictor response and, 793-794

Resource allocation, for geriatric patients, 176

Respiratory depression, from intrathecal analgesia, 388–389

Respiratory function, in elderly, 47–58 alterations in ventilatory drive and control, 53–54

changes in respiratory mechanics and lung volumes, 49–52 clinical and anesthetic implications, 54

impaired efficiency of gas exchange, 52–53

structural alterations in upper and lower airways, 48–49

Respiratory mechanics, changes in, in elderly, 49–52

Respiratory problems, after general anesthesia, in pediatric patients, 616 Respiratory symptoms, in terminally ill

patients, 196–198
Respiratory tract infections, as risk for

Respiratory tract infections, as risk for anesthesia in pediatric patients, 602–603

Ring block of the penis, in pediatrics, use of nerve stimulator, 371

Ropivacaine, for epidural anesthesia, 260, 261

for local anesthesia, 230

in pediatric regional anesthesia, 376 for regional anesthesia for obstetrics, 400

for spinal anesthesia, 236, 242

transient neurologic symptoms from, 238

Sacroiliac joint injection, for chronic pain management, 435–438

Sciatic nerve block, 323, 325 anterior technique, 325–326, 327 classic technique, 325

complications, 326

in pediatrics, 372–373

Scrotum, neuroanatomy of, 887 Sedation, during regional anesthesia, 99 Selective nerve root block, for chronic pain management, 439–441

Sepsis, in pathogenesis of renal insufficiency, 779

as risk factor for acute renal failure, 707 as vasomotor nephropathy, 800

vasopressor therapy in, 801

Septicemia, transient, as complication of transurethral resection of prostate, 893

Serum potassium, preoperative evaluation, for kidney transplant surgery, 924–925 Sickle cell anemia, as risk for anesthesia in pediatric patients, 609–610

Sickle cell disease, complications from, in kidney transplant surgery, 942

Skin damage, as complication of

positioning aged patients, 113–114 Sodium bicarbonate, in cardiopulmonary resuscitation in the elderly, 153

resuscitation in the elderly, 153 with local anesthetics for lower extremity nerve blocks, 337

Somatic nerve blocks, in chronic pain management, blockade of stellate ganglion, 442–445

celiac plexus block, 447-452

cervical facet, median branch blocks, 432, 434–435

epidural steroid injection, 438-439 facet blocks, 430-431

lumbar facet, median branch blocks, 431-432

lumbar sympathetic blockade, 445–447 sacroiliac joint injection, 435–438 selective nerve root block, 439–441 superior hypogastric plexus block, 452–454

sympathetic nerve blocks, 442 Sore throat, as intubation-related

complication, in pediatric patients, 616 Spinal anesthesia, anesthetic distribution, effect on recovery, 245

predictors, 243 strategies, 243, 245

choice of agent, 240–242 bupivacaine, 241 lidocaine, 241

mepivacaine, 242 opioids, 242

procaine, 241 ropivacaine, 242 tetracaine, 241

complications of, 461-485

anterior spinal artery syndrome, 468 choice of needle and risk of post-dural puncture headache, 239–240

incidence and etiology of, 462, 463 local anesthetic toxicity, 465–466 meningitis and spinal abscess, 474–

478, 479–480 nerve injury from needle and catheter placement, 462, 464

neuraxial block in patients with preexisting neurologic disorders, 480–481

neurologic injury and anticoagulation, 238–239

performance of neuraxial block in patients under anesthesia, 481–482 postdural puncture headache, 464-465 spinal hematoma, 468

antiplatelet medications, 473–474 intravenous and subcutaneous standard heparin, 470–471

low molecular weight heparin, 471–472

oral anticoagulants, 472–473 transient neurologic symptoms, 235, 236, 237–238, 466–468

herpes simplex virus and, 478–479 HIV and, 479

indications for, ambulatory surgery, 237 spinal vs. general anesthesia, 236 types of surgery, 236–237 intrathecal adjuncts, 245–246

local anesthetic dose, 242–243, 244 new developments in, 235–249

for transurethral resection of prostate, 888

vs. epidural anesthesia, for cesarean section, 396

Spinal cord damage, from central blocks, 62

Spinal hematoma, as complication of spinal and epidural anesthesia, 468, 470–474

Spinal subarachnoid anesthesia, in pediatrics, 364–366 dosing, 365–366 risks and complications, 366

technique, 365 Spleen, affected by extracorporeal shock

wave lithotripsy, 963
Stents, endovascular aortic, for prevention of renal failure, 733–734

Steroid therapy, as risk for anesthesia in pediatric patients, 609

Stones, renal, treatment of. See Extracorporeal shock wave lithotripsy and percutaneous nephrolithotomy.

Strategy, in business of perioperative medicine, 692–693

Stroke, from CNS dysfunction after anesthesia, 63

Subacute bacterial endocarditis prophylaxis, before surgery, in pediatric patients, 608

Subarachnoid anesthesia, for transurethral resection of prostate, 888

Subarachnoid space, pressure changes in, from combined spinal-epidural technique, 279–281

Succinylcholine-induced myalgia, postoperative, in pediatric patients, 626

Sufentanil, for epidural anesthesia, 260–261 as opioid analgesic, 261

pharmacology of, in elderly patients, 15–16 for spinal anesthesia, 242

Superior hypogastric plexus block, for chronic pain management, 452–454 Supraclavicular block, for upper extremity

regional anesthesia, 309-310

Surgical clinical pathways, in community hospital practice, 542–543

Surrogate endpoints, in outcomes research, 634–635

Sympathetic nerve blocks, for chronic pain management, 442

Sympathetic nervous system, changes in, as effect of cardiovascular and autonomic aging, 38–39

Sympathoadrenal activation,

vasoconstrictor response and, 792-793

Telecardiology, 668–669 Telecorrectional care, 669 Teledermatology, 669 Telemedicine, 663–676

future perioperative medicine practice, 673–674

historical perspectives, 664–666 medicolegal issues, 673

primer, bandwidth, 666–667

education, 670–671 miscellaneous, 670

standards, 667

systems, 667

telecardiology, 668–669

telecorrectional care, 669 teledermatology, 669

telepathology, 668

telepsychiatry, 669 teleradiology, 667–668

telesurgery, 670

regulatory issues, 672 reimbursement, 671-672

Telepathology, 668 Telepsychiatry, 669

Teleradiology, 667–668

Telesurgery, 670

Terminal sedation, 173-174

Tetracaine, for local anesthesia, 225–226 for spinal anesthesia, 241

transient neurologic symptoms from, 238 opental, pharmacology of, in elderly

Thiopental, pharmacology of, in elderly patients, 5–6, 7, 8

Thoracic epidural anesthesia, and myocardial ischemia, 414–418

Thoracic paravertebral block, 352–353 Thromboembolism, with epidural

anesthesia and analgesia, 423 Toxic nephropathy, in pathogenesis of renal insufficiency, 780–781 Tramadol, in palliative care, 191 Transdural drug transport, with combined spinal-epidural technique, 278–279

Transient neurologic symptoms, as complication of spinal and epidural anesthesia, 466–468

from spinal anesthesia, 235, 236, 237–238 Transplant surgery, kidney. See *Kidney*

transplant surgery. liver, as risk factor for acute renal failure, 710

Transurethral resection of the prostate, 883–897

advantages of regional anesthesia over general anesthesia, 888–891

alternatives to, 895

anesthetic techniques, 888

complications, absorption of irrigating solution, 891

ammonia toxicity, 892–893

bleeding and coagulopathy, 894 circulatory overload, hyponatremia,

and hypo-osmolality, 891–892 glycine toxicity, 892

hypothermia, 894

perforation of bladder, 893

transient bacteremia and septicemia, 893

TURP syndrome, 894–895

irrigation solutions, 884-885

neuroanatomy of penis, prostate, and bladder, 885–887 pathophysiology of hypertrophic pros-

tate, 883–884 surgical procedures, 884

Trauma surgery, as risk factor for acute renal failure, 709, 710

Treatment redirection, for geriatric patients, 171–172

Trendelenburg position, disadvantages of, in aged patients, 115–116

Tube feeding, in terminally ill, 202 TURP syndrome, as complication of transurethral resection of prostate,

Ularitide, for prevention of acute renal failure, 713

Ultrafiltration, in dialysis, 854, 857–858 Upper airway obstruction, postoperative, in pediatric patients, 617–618

Upper extremity nerve blocks, 297-317 anatomy and, 297-301, 302

common approaches and techniques for, 307–308

axillary blockade, 311–312 infraclavicular block, 310–311 interscalene block, 308–309

intravenous regional anesthesia, 312–

313

894-895

Upper extremity nerve blocks (Continued) isolated peripheral nerve blocks, 313–314

supraclavicular block, 309–310 decreasing latency and beating the clock, 301, 303–305

future of, 314-315

prolonging, 305–307 Uremia, opioids in patients with, 930 Uremic coagulopathy, in patients with chronic renal failure, 869

Urethra, neuroanatomy of, 885–886 prostatic, neuroanatomy of, 886–887

Urinary tract stone disease, treatment of. See Extracorporeal shock wave lithotripsy and percutaneous nephrolithotomy.

Urodilatin, for renal preservation, 760–761 Uterine perfusion, and regional anesthesia for obstetrics, 397

Vascular surgery, anesthesia-related complications after, 96

regional vs. general anesthesia in, 410–413

Vascular system, affecting positioning of aged patients, 109–112

Vasoconstrictor responses, in normal regulation of renal perfusion, 792–795 Vasodilator responses, in normal

regulation of renal perfusion, 796–798 Vasomotor nephropathy, acute tubular necrosis, 798–799 cardiopulmonary bypass, 803-804 liver failure (hepatorenal syndrome),

802–803

nephrotoxic (drug-induced), angiotensinconverting enzyme inhibitors, 804– 805

cyclosporine A, 805

nonsteroidal anti-inflammatory drugs, 804

renal blood flow regulation, autoregulation, and, 791-807

septic shock, 800

vasopressor therapy in, 801–802 Vasopressin, in cardiopulmonary

Vasopressin, in cardiopulmonary resuscitation in the elderly, 151

Vasopressors, for maternal hypotension from regional anesthesia for obstetrics, 399–400

Vecuronium, pharmacology of, in elderly patients, 24–26

Venous stiffening, as effect of cardiovascular and autonomic aging, 38

Ventilatory drive and control, alterations in, in elderly, 53–54

Verapamil, for renal preservation, 757 Volume loading, for intraoperative prevention of renal failure, 731–732

Vomiting, postoperative, 577 in pediatric patients, 619–621 in terminally ill patients, 195–196

